

The genus *Atanycolus* Foerster (Hymenoptera, Braconidae, Braconinae) in China, with description of one new species

Yi-ping Wang^{1,2,†}, Min Shi^{1,‡}, Xue-xin Chen^{1,§}, Jun-hua He^{1,|}

1 State Key Laboratory of Rice Biology, Institute of Insect Science, College of Agriculture and Biotechnology, Zhejiang University, Hangzhou 310029, China **2** College of Forestry and Biotechnology, Zhejiang Forestry College, Lin'an 311300, China

† [urn:lsid:zoobank.org:author:D8E7A259-C421-4F85-BB02-B78179DBABC5](https://zoobank.org/urn:lsid:zoobank.org:author:D8E7A259-C421-4F85-BB02-B78179DBABC5)

‡ [urn:lsid:zoobank.org:author:5983C9E3-1A63-4B10-B50E-A781F8742E49](https://zoobank.org/urn:lsid:zoobank.org:author:5983C9E3-1A63-4B10-B50E-A781F8742E49)

§ [urn:lsid:zoobank.org:author:0054AC5F-7F6D-4811-A560-90983889C861](https://zoobank.org/urn:lsid:zoobank.org:author:0054AC5F-7F6D-4811-A560-90983889C861)

| [urn:lsid:zoobank.org:author:CDDFF38D9-E9AE-4C8F-99CE-23151D3878F3](https://zoobank.org/urn:lsid:zoobank.org:author:CDDFF38D9-E9AE-4C8F-99CE-23151D3878F3)

Corresponding author: Xue-xin Chen (xxchen@zju.edu.cn)

Academic editor: Kees van Achterberg | Received 18 August 2009 | Accepted 3 September 2009 | Published 9 November 2009

[urn:lsid:zoobank.org:pub:8D1AFFE7-12BA-47F3-B581-CF7FFEAEFF7A](https://zoobank.org/urn:lsid:zoobank.org:pub:8D1AFFE7-12BA-47F3-B581-CF7FFEAEFF7A)

Citation: Wang Y-P, Shi M, Chen X-X, He J-H (2009) The genus *Atanycolus* Foerster (Hymenoptera, Braconidae, Braconinae) in China, with description of one new species. ZooKeys 27: 31–41. doi: 10.3897/zookeys.27.251

Abstract

The species of the genus *Atanycolus* Foerster (Hymenoptera, Braconidae, Braconinae) from China are revised and six species are recognized, including one new species (*A. grandis* Wang & Chen, **sp. n.**), which is described and illustrated in this paper. Three species, i.e. *A. crenulatus* Telenga, 1936, *A. lindemani* Tobias, 1980 and *A. sculpturatus* (Thomson, 1892), are reported from China for the first time. A key to species of this genus is provided.

Keywords

Hymenoptera, Braconidae, Braconinae, *Atanycolus*, new species, new record, Coleoptera, parasitoid, Palearctic region, Oriental region, China

Introduction

Atanycolus Foerster is a cosmopolitan genus with 59 described species (Yu et al. 2007). Most of species are less than 10.0 mm long. They are ectoparasitoids of the larvae of various species of wood and bark-boring beetles (Coleoptera: Buprestidae, Cerambycidae and Scolytidae), many of which are notorious pests of coniferous and broadleaved trees. The chemical control of these pests is difficult because they live under bark. Therefore, the species of this genus may be important natural control agents of many of these beetles.

This is a part of an on-going study of the subfamily Braconinae (Hymenoptera: Braconidae) of China (Wang et al. 2003a, b, c, d, 2004, 2006a, b, c, d, 2007, 2008). In this study, six species of genus *Atanycolus* were found to occur in China, of which one species is new to science (*A. grandis* sp. n.) and three species are new to China (*A. crenulatus* Telenga, 1936, *A. lindemani* Tobias, 1980 and *A. sculpturatus* (Thomson, 1892)). The new species is described and illustrated below. A key to the Chinese species of the genus is provided. The examined specimens are deposited in the Parasitic Hymenoptera Collection, Zhejiang University, Hangzhou, China (ZJUH), Institute of Plant Physiology and Ecology, the Chinese Academy of Sciences, Shanghai (SIPPE) and the Institute of Zoology, the Chinese Academy of Sciences, Beijing (IZBJ).

The morphological terminology used in this paper follows that of van Achterberg (1979), Harris (1979) and Quicke (1987). All descriptions and measurements were made under a Leica MZ 12.5 stereomicroscope (Wetzlar, Germany), and photos taken by a digital camera (Q-Imaging, Micropublisher 3.3 RTV) attached to a Leica MZ APO stereomicroscope (Wetzlar, Germany) using Synoptics Auto-Montage version 5.0 software.

Key to species of genus *Atanycolus* Foerster in China

1. Second metasomal suture (= groove between second and third metasomal tergites) wide and strongly crenulate; third tergite with strongly raised area antero-laterally **2**
- Second metasomal suture narrow and weakly crenulate medially, and at least laterally smooth; third tergite with weakly raised area antero-laterally **5**
2. Head entirely blackish; longitudinal grooves of second metasomal tergite shallow laterally; first metasomal tergite blackish brown ***A. denigrator* (Linnaeus, 1758)**
- Head dark yellow or yellowish red; longitudinal grooves of second metasomal tergite deep laterally; first metasomal tergite black or yellow **3**
3. Triangular area of third and fourth metasomal tergites (except for medio-basally) uniformly sculptured; metasoma pale yellow and head blackish brown ***A. sculpturatus* (Thomson, 1892)**
- Third and fourth tergites smooth without any sculpture; metasoma yellowish brown or blackish brown and head reddish yellow **4**

4. Length of body more than 10 mm; ocellar triangular area with blackish spot (Fig. 2); wing membrane infuscated (Fig. 5); third metasomal tergite with strongly raised area antero-laterally (Fig. 7, 8)..... ***A. grandis* Wang & Chen, sp. n.**
- Length of body less than 10 mm; ocellar triangular area reddish yellow; wing membrane yellowish brown; third tergite with weakly raised area antero-laterally..... ***A. crenulatus* Telenga, 1936**
5. Second metasomal tergite with parallel longitudinal impressed grooves laterally, medio-basal smooth triangular area deeply impressed; suture between second and third tergites crenulate medially and smooth laterally; tergites three to six orangish yellow ***A. initiator* (Fabricius), 1793**
- Second tergite without parallel longitudinal impressed grooves laterally, medio-basal smooth triangular area without a deep impression; suture between second and third tergites completely smooth; tergites three to six yellowish-white..... ***A. lindemani* Tobias, 1980**

Descriptions

Atanycolus grandis Wang & Chen, sp. n.

urn:lsid:zoobank.org:act:6AA69564-8B04-4CF7-BACE-1C3F865B6C95


Figs 1–9

Length of body 10.0–14.0 mm, fore wing 8.0–12.0 mm, ovipositor sheath 10.0–12.5 mm.

Head. Antenna with 68 segments; terminal flagellomere tapering apically, approximately 1.4 times longer than basal width; first flagellomere parallel-sided but weakly flared basally, 1.2 and 1.5 times longer than the second and third ones, respectively; the latter 1.4 times longer than its maximum width; median flagellomeres as long as its maximum width; transverse medial clypeal carina with a row sparse long setae; height of clypeus: inter-tentorial distance: tentorio-ocellar distance = 5: 9: 6; face with sparse long setae, width of face: width of head: maximum length of eye in dorsal view = 18: 34: 19; frons strongly impressed medially; shortest distance between posterior ocelli: diameter of posterior ocellus: shortest distance between posterior ocellus and eye = 2.5: 2.0: 9; vertex smooth and shiny with dense setae.

Mesosoma. Length of mesosoma 1.8 times as long as its maximum height, smooth and shiny with sparse long setae; notauli deeply impressed anteriorly and shallowly posteriorly, with sparse short setae along entire length; scutellar sulcus relatively narrow and deep, and distinctly crenulate; metanotum with strongly raised area medially; propodeum smooth, with sparse setae, but relatively densely and long setose laterally.

Wing. Length of fore wing veins SR1: 3-SR: r = 64: 45: 12; vein 1-SR+M slightly bent basally; length of fore wing veins 2-SR: 3-SR: r-m = 21: 45: 19; vein cu-a interstitial. Hind wing: length of veins SC+R1: 2-SC+R: 1r-m = 30: 6: 12; vein C+SC+R with three thickened (humular) bristles apically.



Figures 1–9. *Atanycolus grandis* Wang & Chen, sp. n.: **1** head, frontal view **2** head, dorsal view **3** scapus, lateral view **4** scapus, ventral-lateral view **5** wings **6** metanotum and propodeum, dorsal view **7** metasomal tergites, dorsal view **8** metasomal tergites, lateral view **9** apex of ovipositor, lateral view.

Leg. Length of fore femur: tibia: tarsus = 50: 53: 94; length of hind femur: tibia: basitarsus = 15: 25: 9, 3.2, 10.0 and 6.1 times their maximum width, respectively; tibia of hind leg with longitudinal groove medially.

Metasoma. First metasomal tergite 1.2 times longer than its maximum apical width, strong apical dorsal carinae occupying basal 3/4 of its length; second tergite with a glabrous medio-basal triangular area and sublateral longitudinal grooves, the remainder deeply striate; third tergite with distinct carina antero-laterally, smooth and shiny medially, distinctly sculptured postero-laterally; suture between second and third metasomal tergites wide, deep and crenulate; third tergite with transverse deeply impressed groove apically; tergites four to six smooth and shiny;

hypopygium acute apically, slightly extending beyond apex of metasoma; ovipositor sheath as long as fore wing, and ovipositor with ventral teeth sub-apically.

Colour. Antenna blackish; head dark yellow except for ocellar triangular area with blackish spot and maxillary palp reddish-yellow; mesosoma reddish-yellow but mesoscutum with blackish-brown spots laterally; propodeum dark brown; fore leg reddish-yellow, middle and hind legs blackish-brown; wings membrane infuscated and veins blackish brown; metasomal tergites black; ovipositor sheath blackish-brown.

Variation. Paratypes similar to the holotype. Antenna with 66–69 segments. Head in dorsal view 1.6–1.7 times as broad as long. Hind femur 3.1–3.3 times as long as broad medially. Pterostigma somewhat wide, 3.1–3.2 times as long as wide; second submarginal cell of fore wing long, vein 3-SR 2.2–2.3 times as long as the 2-SR. First tergite 1.1 times as long as broad posteriorly. Second tergite distinctly longer than third tergite. Vertex pale yellow or reddish yellow; mesoscutum with blackish brown spots laterally; middle and hind legs pale yellow; ovipositor sheath yellowish brown.

Specimens examined. Holotype, ♀, West Tianmu Mountain (119°23'E, 30°18'N), Zhejiang, 2000.VII.3, Ma Yun, No. 200103596 (ZJUH). Paratypes: 1♀, T'ienmu-Shan (119°23'E, 30°18'N), China, 1937.VII.2, O. Piel., Musee Heude (SIPPE); 1♀, T'ienmu-Shan (119°23'E, 30°18'N), China, 1936.VII.6, O. Piel., Musee Heude (SIPPE); 4♀♀, Songyang (119°29'E, 28°28'N), Zhejiang, China, 1989.VIII.16, He Jun-hua, 1989.VII.11–14, 11–21, No. 895251, 895313, 895448, 895251 (ZJUH); 2♀♀, Zhuchaoguan (119°30'E, 28°29'N), Songyang, Zhejiang, China, 1994.VII.17, Cai Ping, No. 944170, 944172 (ZJUH); 1♀, Zhuchaoguan (119°30'E, 28°29'N), Songyang, Zhejiang, China, 1994.VII.17, Xu Zai-fu, No. 944369 (ZJUH); 1♀, Baishanzu (119°14'E, 27°52'N), Qingyuan, Zhejiang, China, 1993.VII.30, Wu Hong, No. 945161 (ZJUH); 1♀, Wuyi Mountain (118°02'E, 27°46'N), Fujian, China, 1986.VI.16, Wang Jia-she, No. 870644 (ZJUH); 1♀, Huaping (110°2'E, 25°10'N), Longsheng, Guangxi, China, 1982. VI. 25, He Jun-hua, No. 823297 (ZJUH); 1♀, Longgang (108°23'E, 34°19'N), Longzhou, Guangxi, China, 1982.V.20, He Jun-hua, No. 821594 (ZJUH); 1♀, Ziping (114°16'E, 26°56'N), Jinggang mountain, Jiangxi, China, 1981.V.28, Liu Jin, Liu Yao, No. 340041551 (ZJUH); 1♀, Pingba (106°27'E, 26°42'N), Guizhou, China, 19??, Guizhou Forest Academy Institute, No. 801723 (ZJUH).

Biology. Unknown.

Etymology. From latin “grandis” meaning for large, referring to large body of type specimens.

Diagnosis. This species is similar to *Atanycolus crenulatus* Telenga, but can be separated from the latter by having the length of body more than 10 mm; the wings infuscated (Fig. 5); ocellar triangular area with a blackish spot (Fig. 2); vein 1-SR+M of fore wing slightly bent basally (Fig. 5); propodeum with relatively dense and long setae laterally (Fig. 6); second tergite deeply striate except for smooth medio-basal triangular area (Fig. 7), and third metasomal tergite with a distinctly raised area antero-laterally (Figs 7, 8).

***Atanycolus crenulatus* Telenga, 1936**

Atanycolus crenulatus Telenga, 1936: 327; Telenga, 1952: 120 (Trans. 1969: 95); Tobias, 1971, 54: 207; Tobias et al., 2000, 4: 183.

Specimens examined: CHINA: GANSU: 1♀, Lanzhou (103°44'E, 36°2'N), 1980. VII.29, Wang Chang-zhong, No. 853597 (ZJUH).

Biology. unknown.

Distribution. China (Gansu) and Russia.

Note. This species is new to China.

***Atanycolus denigrator* (Linnaeus, 1758)**

Ichneumon denigrator Linnaeus, 1758, 10a 1: 563.

Bracon denigrator: Panzer, 1801: 163.

Atanycolus denigrator: Foerster, 1862, 19: 238; Shenefelt, 1978: 1441; Papp, 1998: 147; Sheng, 1990, 12 (1): 56.

Biology. Reported from the following hosts: *Acanthocinus aedilis* L., *Rhagium indagator* F., *R. inquisitor* L., *Saperda populnea* L. and *Tetropium fuscum* L. (Coleoptera: Cerambycidae), *Anthaxia morio* F. and *Chrysobothris chrysostigma* L. (Coleoptera: Buprestidae).

Distribution. China (Inner Mongolia); Austria; Bulgaria; Croatia; Czechoslovakia; Finland; France; Germany; Greece; Hungary; Israel; Italy; Kazakhstan; Korea; Mongolia; Niger; Norway; Poland; Russia; Slovakia; Sweden; Switzerland; Turkey; United Kingdom.

Note. This species has been reported from Daxing'an Mountain (China) by Sheng (1990), but no specimens were available for our study.

***Atanycolus initiator* (Fabricius, 1793)**

Ichneumon initiator Fabricius, 1793, 2: 161.

Bracon (Coeloides) initiator: Blanchard, 1840, 3: 341.

Bracon (Vipio, Coelobracon) genalis Thomson, 1892, 17: 1800. syn. by Roman, 1912.

Atanycolus initiator: Szepliget, 1901, 33: 176; Watanabe, 1950, 21(2): 20 (Shanxi of China); Papp, 1998a, 59: 147; Tobias et al., 2000, 4: 183; Sheng, 1990, 1: 33.

Atanycolus mongolicus Telenga, 1936, 5 (2): 92. syn. by Tobias, 1971.

Atanycolus petiolaris Thomson, 1892, 17: 1859. syn. by Tobias, 1986.

Specimens examined. CHINA: HEILONGJIANG: 6♀♀, Dailing (128°54'E, 47°40'N), Yichun, 1956.VI.10, Shi Zhen-hua, No. 5710. 3, Ex. *Monochamus sutor* (L.); 20♀♀, Dailing (128°54'E, 47°40'N), Yichun, 1977.VII.24, He Jun-hua, No. 771811, 771749, 771771, 771786, 771845, 771870, 771680, 771703, 771713, 771710, 771850, 771802, 771887, 771892, 771812, 771687, 771725, 771813, 771726, 771727, 771903 (ZJUH); CHINA: INNER MONGOLIA: 3♀♀2♂♂, Xiao Xing'anling (127°42'E, 46°28'N), 1955.VI.1, Collector unknown, No. 5607. 1, Ex. *Ips typographus*, L. (ZJUH); CHINA: SHANXI: 1♀1♂, Taiyuan (112°33'E, 37°54'N), 1987.V, Zhao Rui-liang, No. 870074, 870075 (ZJUH); CHINA: SHANGDONG: 1♀, Tai mountain (117°59'E, 36°28'N), Tai'an, 1997.VI.17, Li Qiang, No. 200011169 (ZJUH); CHINA: HUBEI: 2♀♀, Jingshan (113°23'E, 31°6'N), 1980.VI, Zhan Zhong-cai, No. 824303, 824302 (ZJUH).

Biology. The known hosts are *Monochamus sutor* (L.) (Coleoptera: Cerambycidae) and *Ips typographus* L. (Coleoptera: Scolytidae) according to collecting labels. Other reported hosts are *Semanotus rufipennis* Motschulsky, *Acanthocinus aedilis* L., *Cerambyx scopolii* Fuessly, *Criocephalus rusticus* L., *Rhagium inquisitor* L., *Tetropium castaneum* L. and *T. fuscum* Fabricius (Coleoptera: Cerambycidae), *Aegeria flaviventris* Staudinger and *A. vespiformis* L. belonging to family Sesiidae of Lepidoptera based on the catalogues of Shenefelt (1978) and Yu et al. (2005).

Distribution. China (Heilongjiang, Inner Mongolia, Shanxi, Shandong, Henan and Hubei); Austria; Azerbaijan; Croatia; Czech Republic; Czechoslovakia; Finland; France; Germany; Hungary; Italy; Japan; Latvia; Lithuania; Mongolia; Norway; Poland; Russia; Slovakia; Spain; Sweden; Switzerland; Turkey; Turkmenistan; Ukraine; United Kingdom.

Atanycolus lindemani Tobias, 1980

Atanycolus lindemani Tobias, 1980, 7: 289–295; Tobias and Belokobylskij, 2000, 4: 184.

Specimens examined. CHINA: HEILONGJIANG: 1♂, Dailing (128°54'E, 47°40'N), Yichun, 1956.V.29, Shi Zhen-hua, No. 5710.1 (ZJUH); CHINA: HEILONGJIANG: 1♂, Dailing (128°54'E, 47°40'N), Yichun, 1977.VII.24, He Jun-hua, No. 771808 (ZJUH); CHINA: JILIN: 1♀, Gongzhu Mountain (124°49'E, 43°3'N), 1983.VIII.9, Wang Cheng-lun, No. 840129 (ZJUH); CHINA: XINJIANG: 1♀1♂, Shihezi (86°0'E, 44°18'N), 1981.VI.15, He Fu-de, No. 816488, Ex. *Scolytus seulensis* Murayama (ZJUH); CHINA: SHAANXI: 1♀1♂, Dingbian (107°59'E, 37°60'N), 1981.V.23, Dang Xin-de, No. 815987, Ex. *Xyloterinus politus* Say (ZJUH).

Biology. The known hosts are *Xyloterinus politus* Say and *Scolytus seulensis* Murayama (Coleoptera: Scolytidae) according to collecting labels.

Distribution. China (Heilongjiang, Jilin, Xinjiang and Shaanxi) and Russia.

Note. This species is new to China.

Atanycolus sculpturatus (Thomson, 1892)

Bracon (*Vipio*, *Coelobracon*) *sculpturatus* Thomson, 1892, 17: 1800.

Vipio (*Coelobracon*) *sculpturatus*: Kriechbaumer, 1898, 24: 246.

Coelobracon sculpturatus: Szepligeti, 1896, 19: 168.

Coeloides (*Atanycolus*) *sculpturatus*: Marshall, 1897, 5 bis: 118.

Vipio (*Atanycolus*) *sculpturatus*: Schmiedeknecht, 1897 1: 512.

Atanycolus sculpturatus: Dalla Torre, 1898, 4: 296; Papp, 1998b, 59: 147; Tobias et al., 2000, 4: 183.

Iphiaulax (*Atanycolus*) *sculpturatus*: Hellen, 1927, 56: 11.

Atanycolus signatus Szepligeti, 1901, 33: 176. syn. by Papp, 1960.

Coelobracon signatus: Fahringer, 1926, 1(2–3): 136.

Specimens examined. CHINA: XINJIANG: 1♀, Liudaowan (87°65'E, 43°74'N), 1982.VII.28, No. 860047, collector unknown; 1♀, Shihezi (86°00'E, 44°18'N), 1981.VIII.14, He Fu-de, No. 820018 (ZJUH).

Biology. According to Shenefelt (1978) Yu et al. (2007), the known hosts are *Chrysobothris sorieri* Lap. and *Agrilus biguttatus* F. (Coleoptera: Buprestidae), *Tetropium gabrieli* Weise, *Ruguloscolytus mediterraneus* Egg. and *Leptura rubra* (L.) (Coleoptera: Cerambycidae).

Distribution. China (Xinjiang); Austria; Croatia; Czech Republic; Czechoslovakia; Finland; France; Germany; Greece; Hungary; Italy; Japan; Russia; Slovakia; Switzerland; Ukraine.

Note. This species is new to China.

Acknowledgements

We thank Dr. C. van Achterberg (Leiden, The Netherlands) for his critical review of an early version of the manuscript and for improving the text. We also thank two anonymous reviewers for their suggestions and improvement of the text. The project was jointly supported by funds from the National Science Fund for Distinguished Yuong Scholars (No. 30625006), the National Natural Science Foundation of China (Nos. 30700063, 30499341), the National S&T Infrastructure Project (Nos. 2005DKA21402, 2005DKA21105), and the National Special Basic Research Funds (Nos.2006FY110500-3, 2006FY120100).

References

- Achterberg C van (1979) A revision of the subfamily Zelinae auct (Hymenoptera: Braconidae). *Tijdschrift voor Entomologie* 122: 241–479.
- Ashmead WH (1900) Classification of the ichneumon flies, or the superfamily Ichneumonoidea. *Proceedings of the United States National Museum* 23: 1–120.
- Blanchard E (1840) *Historie naturelle des insects, 3. Orthoptera, Neuropteres, Hemipteres, Hymenopteres, Lepidopteres, et Dipteres, avec une introduction rendrant L'anatomie et la physiologie des Animaux Articulés par M. Brull, Paris, 672 pp.*
- Dalla Torre CG (1898) *Catalogus Hymenopterorum, 4, Braconidae. G. Engelmann, Leipzig, 323 pp.*
- Fabricius W (1793) *Entomologia Systematica. Hafniae, C.G. Proft. 2, 517 pp.*
- Fahringer J (1928–35) *Opuscula braconologica. 2. Aethiopische Region, Lieferung 1–8, 635 pp.*
- Foerster A (1862) Synopsis der Familien und Gattungen der Braconen. *Verhandlungen des naturhistorischen Vereins de Preussischen Rheinlande und Westfalens* 19: 225–288.
- Harris RA (1979) A glossary of surface sculpturing. *Occasional Papers of the Bureau of Entomology of the California Department of Agriculture* 28: 1–31.
- Hellen W (1927) Zur Kenntniss der Braconiden (Hym.) Finnlands. I. Subfam. Braconinae (part.), Rhogadinae und Spathiinae. *Acta Societatis pro Fauna et Flora Fennica* 56(12): 1–59.
- Linnaeus C von (1758) *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species cum characteribus, differentiis, synonymis locis. Tomus I. Editio decima, reformata. Laurnetii Salvii, Holmiae, 824 pp.*
- Marshall TA (1897) Les Braconides. In: Andre E (ed). 1897–1900. *Species des Hymenoptera d'Europe et d'Algerie* 5: 118.
- Panzer GWF (1801) *Systematische Nomenclaturganer weiland Herrn Dr. Jacob Christian Schonfers natanlich ausgemahlte Abbildungen regensburgischer Insekten. Teil I-III. Johann Jakob Palm, Erlangae, 260 pp.*
- Papp J (1960) A revision of the tribe Braconini Ashmead from the Carpathian Basin (Hymenoptera, Braconidae). *Annales Historico-Naturales Musei Nationalis Hungarici* 52: 367–380.
- Papp J (1998a) Contribution to the braconid fauna of the former Yugoslavia, 4. Braconinae (Hymenoptera: Braconidae). *Folia Entomologica Hungarici* 59: 145–162.
- Papp J (1998b) New braconid wasps (Hymenoptera, Braconidae) in the Hungarian Natural History Museum, 6. *Annales Historico-Naturales Musei Nationalis Hungarici* 90: 221–256.
- Quicke DLJ (1987) The Old World genera of braconine wasps (Hymenoptera: Braconidae). *Journal of Natural History* 21: 43–157.
- Quicke DLJ, Sharkey MJA (1989) A key to and notes on the genera of Braconinae (Hym. Braconidae) from America north of Mexico with descriptions of two new genera and three new species. *The Canadian Entomologist* 121: 337–361.
- Schmiedeknecht O (1897) Das Studium der Braconiden nebst einer revision der europäischen und benachbarten Arten der Gattungen *Vipio* and *Bracon*. *Illustrierte Wochenschrift Entomologie* 1: 496–592.

- Shenefelt RD (1978) Hymenopterorum catalogus (nov. editio). Pars 15. Braconidae 10. Braconinae, Gnathobraconinae, Mesostoinae, Pseudodicrogeniinae, Telengainae, Ypsistocerinae plus Braconidae in general, major groups, unplaced genera and species. Dr. W. Junk B.V., The Hague, Holland, 1425–1872.
- Sheng ML (1990) New records of Hymenoptera parasitizing wood-boring insects from China. *Entomotaxonomia* 12(1): 56.
- Sheng ML, Liu ZF, Cui ZQ (1992) Natural enemies of the tree borers in fire-slash in Daxing'anling mountains. *Scientia Silvae Sinicae* 28 (1): 80–84.
- Szépligeti G (1896) Adatok a magyar fauna braconidainak ismeretehez. *Beiträge zur Kenntnis der ungarischen Braconiden. Termeszetráji Füzetek* 19: 165–186.
- Szépligeti G (1901) Tropische Cenocoelioniden und Braconiden aus der Sammlung des Ungarischen National-Museums. *Termeszetráji Füzetek* 24: 353–402.
- Telenga NA (1936) Faune de l'U. S. S. R. Insectes Hymenopteres. Fam. Braconidae (P.I.). *Inst. zool. Akad. Nauk SSSR, Leningrad, Fauna Rossii* 5(2): 84–402.
- Telenga NA (1952) The origin and evolution of parasitism in parasitic Hymenoptera and the formation of their fauna in the U.S.S.R. *Akademia nauk U.S.S.R., Kiev*, 137 pp.
- Thomson CG (1892) XLIV, Bidrag till Braconidernas Kannedom. *Opuscula Entomologica* 16: 1659–1751.
- Tobias VI (1971) Review of the Braconidae (Hymenoptera) of the USSR. *Trudy Vsesoyuznogo Entomologicheskogo Obshchestva* 54: 156–269, 962–964.
- Tobias VI (1980) New species of braconids (Hymenoptera, Braconidae) - parasites of bark beetles from Mongolia and the USSR. *Nasekomye Mongolii* 7: 289–295.
- Tobias VI, Belokobylskij AB (2000) Key to the insects of Russian Far East Vol. VI. Newropteroidea. Mecoptera. Hym. Pt. 4. Vladivostok: Dal nauka., 649 pp.
- Viereck HL (1912) Contributions to our knowledge of bees and ichneumon-flies, including descriptions of twenty-one new genera and fifty-seven new species of ichneumon-flies. *Proceedings of the United States National Museum* 42: 613–648.
- Wang YP, Chen XX, He JH (2003a) Taxonomic studies on the subtribe Aphrastobraconina Ashmead (Hymenoptera: Braconidae, Braconinae) in China. *Entomologica Fennica* 14: 118–124.
- Wang YP, Chen XX, He JH (2003b) First Chinese record of the genus *Trispinaria* Quicke (Hymenoptera: Braconidae, Braconinae) and a description of a new species. *Acta Zootaxonomica Sinica* 28: 333–335.
- Wang YP, Chen XX, He JH (2003c) The genus *Testudobracon* Quicke in eastern China and the descriptions of three new species (Hymenoptera: Braconidae: Braconinae). *European Journal of Entomology* 100: 411–416.
- Wang YP, Chen XX, He JH (2003d) The discovery of the genus *Shelfordia* Cameron (Hymenoptera: Braconidae, Braconinae) in China, with description of one new species. *Entomologia Sinica* 10: 215–220.
- Wang YP, Chen XX, He JH (2004) A review of *Bracon* (*Rostrobracon*) (Hymenoptera: Braconidae: Braconinae) from China, with description of new species. *Oriental Insects* 38: 341–346.
- Wang YP, Chen XX, He JH (2006a) The discovery of genus *Spinadesha* Quicke of tribe Adeshini van Achterberg (Hymenoptera: Braconidae: Braconinae) in China, with description of one new species. *Biologia* 61: 145–147.

- Wang YP, Chen XX, He JH (2006b) Three genera with eyes setose of Braconinae (Hymenoptera: Braconidae) in China, with the description of one new species. *Entomotaxonomia* 28: 131–138.
- Wang YP, Chen XX, He JH (2006c) The genus *Pseudoshirakia* van Achterberg (Hymenoptera: Braconidae) from China. *The Pan-Pacific Entomologist* 81: 46–49.
- Wang YP, Chen XX, Wu H, He JH (2006d). The genus *Coeloides* Wesmael of subfamily Braconinae (Hymenoptera: Braconidae) in China. *Zootaxa* 1239: 1–17.
- Wang YP, Chen XX & He JH (2007) The genera *Aspidobracon* van Achterberg and *Philomacroplea* Cameron (Hymenoptera: Braconidae: Braconinae) in China, with descriptions of two new species. *Annals of the Entomological Society of America* 100(3): 390–393.
- Wang YP, Shi M, He JH, Chen XX (2008) Taxonomic study of the tribe Iphiaulacini (Hymenoptera: Braconidae: Braconinae) from China with descriptions of two new species. *Entomotaxonomia* 30(3): 181–195.
- Watanabe C (1950) A preliminary revision of the genus *Aphrastobracon* Ashmead (Hymenoptera, Braconidae). *Journal of the Faculty of Agriculture, Hokkaido (Imp.) University* 48: 291–304.
- Yu DS, van Achterberg C, Horstmann K (2007) *World Ichneumonoidea 2006: Taxonomy, Biology, Morphology and Distribution*. CD/DVD. Taxapad, Vancouver, Canada.